

Photon localization barrier can be overcome

P. Saari^{1,2}, M. Menert¹, and H. Valtna¹

¹*Department of Physics, University of Tartu;* ²*Institute of Physics, University of Tartu; Riia 142, Tartu 51014, Estonia*

Abstract

In contradistinction to a widespread belief that the spatial localization of photons is restricted by a power-law falloff of the photon energy density, I.Bialynicki-Birula [Phys. Rev. Lett. **80**, 5247 (1998)] has proved that any stronger – up to an almost exponential – falloff is allowed. We are showing that for certain specifically designed cylindrical one-photon states the localization is even better in lateral directions. If the photon state is built from the so-called focus wave mode, the falloff in the waist cross-section plane turns out to be quadratically exponential (Gaussian) and such strong localization persists in the course of propagation.

While quantum electrodynamics (QED) underwent an impressive development and reached its maturity in the middle of the last century, one of its basic concepts – the photon wave function in free space – was deprived of such fortune. Although the photon wave function in coordinate representation was introduced already in 1930 by Landau and Peierls [1] the concept was found to suffer from inherent difficulties that were not overcome during the century (see review [2]). The common explanation presented in textbooks (e.g., [3],[4]) may be summed up as follows: (i) no position operator exists for the photon, (ii) while the position wave function may be localized near a space-time point, the measurable quantities like the electromagnetic field vectors, energy, and the photodetection probability remain spread out due to their non-local relation with the position wave function. However, just before the turn of the century both of these widely-espoused notions were disproved [5],[6] and in the new century a fresh interest in the photon localization problem seems to have been awakened (see, e.g., [7],[8],[9]), meeting the needs of developments in near-field optics, cavity QED, and quantum computing.

I. Bialynicki-Birula writes [6] that the statement “even when the position wave function is strongly concentrated near the origin, the energy wave function is spread out over space asymptotically like $r^{-7/2}$ ” (citation from [4], p. 638) is incorrect and that both wave functions may be strongly concentrated near the origin. He demonstrates, on one hand, that photons can be essentially better localized in space – with an exponential falloff of the photon energy density and the photodetection rates. On the other hand, he establishes – and it is even somewhat startling that nobody has done it earlier – that certain localization restrictions arise out of a mathematical property of the positive frequency solutions which therefore are of a universal character and apply not only to photon states but hold for all particles. More specifically, it has been proven in the Letter [6] for the case of spherically imploding-exploding one-photon wavepacket that the Paley-Wiener theorem allows even at instants of maximal localization only such asymptotic decrease of the modulus of the wave function with the radial distance

r that is *slower* than the linear exponential one, i.e., anything slower than $\sim \exp(-Ar)$, where A is a constant. The latter is what the Paley-Wiener theorem says about a function whose Fourier spectrum contains no negative frequencies.

The purpose of the present Letter is to indicate that one-photon wave functions of a specific type can break the localization restriction and exhibit the linear exponential and even faster falloff with the distance. Yet, there is no contradiction either with the result of Ref. [6] or with the Paley-Wiener theorem, since the wave functions are cylindrical and exhibit an exceptionally strong localization in two dimensions out of three.

As an introduction, we consider briefly the simplest case of a one-dimensional Landau-Peierls wave function in order to indicate how the Paley-Wiener theorem restricts the spatial localization of a photon. Then we study the radial falloff for three different cylindrical wave functions, using exactly the same formalism that has been presented in Ref. [6]. Finally, the discussion of our results allows us to refine the analysis given in Ref. [6].

Let us consider a one-photon (1ph) state that corresponds to a plane-wave pulse propagating unidirectionally, say, along the axis z , being polarized along a lateral axis (say, the x axis)

$$|1ph\rangle = \int_0^\infty dk_z f(k_z) a^+(k_z) |vac\rangle, \quad (1)$$

where $a^+(k_z)$ is the creation operator of a photon and $f(k_z)$ is a properly-normalized photon wave function in the momentum representation. Then the inverse Fourier transform (but including positive frequencies only!)

$$\begin{aligned} \Phi(z, t) &= \frac{1}{2\pi} \int_0^\infty dk_z f(k_z) e^{i(k_z z - \omega t)} \\ &= \frac{1}{2\pi} \int_0^\infty dk f(k) e^{ik(z - ct)} \end{aligned} \quad (2)$$

represents the corresponding position space wave function of the photon in state $|1ph\rangle$ (see, e.g., [4], p. 636). The modulus squared $|\Phi(z, t)|^2$ gives the photon probability density, i.e., the degree of localization along the axis

z (in the given case in the directions x and y any localization is absent). If $f(k)$ differs from zero within a wide frequency band, the probability $|\Phi(z, t)|^2$ may be strongly localized around a point z_0 moving along the axis z with the speed of light c . However, since $\Phi(z, t)$, due to the absence of negative frequencies in the integral of Eq.(2), is nothing but a complex analytic signal, according to the Paley-Wiener theorem (or criterion) the asymptotic decrease of $|\Phi(z, t)|^2$ with the distance $r = |z - z_0|$ has to be weaker than $\sim \exp(-Ar)$, where A is a constant. All the more excluded are any finite-support functions in the role of $\Phi(z, t)$. To conclude the introduction, let us notice that if the counterpropagating (with $k_z < 0$) Fourier components are involved in Eq.(2), the Paley-Wiener theorem does not apply at the instant $t = 0$.

Following Ref. [6], we shall study the photon localization by examining the asymptotic behavior of the positive frequency part of the Riemann-Silberstein vector $\mathbf{F}(\mathbf{r}, t)$ (called the energy wave function) which directly determines the energy density of a one-photon state and is conveniently expressed through a "superpotential" $\mathbf{Z}(\mathbf{r}, \tau)$

$$\mathbf{F}(\mathbf{r}, t) = \nabla \times \left[i \frac{\partial}{\partial \tau} \mathbf{Z}(\mathbf{r}, \tau) + \nabla \times \mathbf{Z}(\mathbf{r}, \tau) \right], \quad (4)$$

where $\tau \equiv ct$. The vector field $\mathbf{Z}(\mathbf{r}, \tau)$ is nothing but an analytic signal version of the Hertz potential, i.e., $\mathbf{Z}(\mathbf{r}, \tau) = \mathbf{m}\Psi(\mathbf{r}, \tau)$, where $\mathbf{m}$ is a constant vector that includes the proper normalization factor and $\Psi(\mathbf{r}, \tau)$ is any solution of the scalar wave equation, which is taken in the form of the analytic signal.

As the first example leading to a stronger localization that one might expect from the Paley-Wiener theorem, let us consider the photon field where $\mathbf{m}$ is directed along the axis z (any other orientation gives similar results) and $\Psi(\mathbf{r}, \tau)$ is a superposition of cylindrically symmetric Bessel functions J_0 as a wavepacket with the exponential spectrum and a specific dispersion law for the axial wavenumber $k_z(\omega) = \text{const} = k_0$

$$\Psi(\rho, z, \tau) = \int_{|k_0|}^{\infty} dk J_0(k_\rho \rho) e^{-k\Delta} e^{-i(k\tau - k_0 z)}, \quad (5)$$

where the radial coordinate ρ has been introduced and $k_\rho = (k^2 - k_0^2)^{1/2}$ is the lateral component of the wave vector of the monochromatic plane-wave constituents represented with the weight function $e^{-k\Delta}$ whose width is Δ^{-1} . The integral can be taken with the help of a Laplace transform table and we obtain

$$\mathbf{Z}(\rho, z, \tau) = \mathbf{m} \frac{\exp\left(-|k_0| \sqrt{\rho^2 + (\Delta + i\tau)^2}\right)}{\sqrt{\rho^2 + (\Delta + i\tau)^2}} e^{ik_0 z}. \quad (6)$$

Eq.(6) describes a simple cylindrical pulse modulated harmonically in the axial direction and radially converging (when $\tau < 0$) to the axis and thereafter (when $\tau > 0$)

expanding from it, the intensity distribution resembling an infinitely long tube coaxial with the z axis and with a time-dependent diameter (see Fig. 6. in Ref. [10]). It follows from Eqs. (6) and (4) that

$$|\mathbf{Z}(\rho \rightarrow \infty, z, \tau = 0)| \sim \rho^{-1} \exp(-\rho/l), \quad (7)$$

$$|\mathbf{F}(\rho \rightarrow \infty, z, \tau = 0)|^2 \sim [\rho^{-2} + O(\rho^{-3})] \exp(-2\rho/l), \quad (8)$$

where $l \equiv |k_0|^{-1}$ is the characteristic length (or length unit). Thus, while the photon is delocalized in the axial direction, its energy density falloff in the lateral directions is exactly the linear exponential one at all times the conditions $\tau \ll \rho \gg \Delta$ are fulfilled, see Fig. 1. The time derivative as well as the spatial derivatives contain the same exponential factor, ensuring the exponential falloff of the the Riemann-Silberstein vector in Eq.(8). Hence,




FIG. 1: Curves of the radial dependence in a decimal logarithmic scale. Curve A is for $|\mathbf{Z}(\rho, 0, \tau = 0)|$; B, $|\mathbf{Z}(\rho, 0, \tau = 2.5l)|$; C is the same as B but with Ψ taken from Eq. (9); D, $|\frac{\partial}{\partial \tau} \mathbf{Z}(\rho, 0, \tau = 0)|$; E is a reference curve $\exp(-\rho/l)$. The curves A, B, and C have been normalized so that $|\mathbf{Z}(0, 0, 0)| = 1$. The values of the remaining free parameters are $\Delta = 0.1l$ and $\beta = 0.8$.

a one-photon field given by Eq.(6) serves as the first and simplest example where the localization in two transversal dimensions is governed by different rules than localization in three dimensions according to Ref. [6].

The next example is readily available via the Lorentz transformation of the wave function given by Eq.(5) along the axis z , which gives another possible solution of the scalar wave equation. The result is a new independent solution but it can also be considered as the wave given by Eqs. (5) and (6), which is observed in another inertial

reference frame [10]:

$$\Psi(\rho, z, \tau) = \frac{\exp\left(-|k_0| \sqrt{\rho^2 + (\Delta - i\gamma(\beta z - \tau))^2}\right)}{\sqrt{\rho^2 + (\Delta - i\gamma(\beta z - \tau))^2}} \times \exp(i\gamma k_0(z - \beta\tau)), \quad (9)$$

where the relativistic factors $\gamma \equiv (1 - \beta^2)^{-1/2}$ and $\beta \equiv v/c < 1$ have been introduced, v being a free parameter – the relative speed between the frames. In the waist region (see Fig. 2) this wave function has the same radial falloff as was given by Eq.(7), see curve "C" in Fig. 1, while the axial localization follows a power law. The strongly localized waist and the whole amplitude distribution move rigidly and without any spread along the axis z with a superluminal speed c/β . Such wave with intriguing properties, named the focused X wave (FXW) [11], belongs to the so-called propagation-invariant localized solutions to the wave equation – a research field emerged in the 1980-ies (see reviews [11]-[15]) and recently reached its first experimental results [16]-[21]. Hence, in its waist




FIG. 2: The superluminal FXW given by Eq. (9). Shown are the dependences (a) of the modulus and (b) of the real part of the wavefunction on the longitudinal (z , increasing to the right) and a transverse (say, x) coordinates. The distance between the grid lines on the basal plane (x, z) is 22λ , where $\lambda = 2\pi|k_0|^{-1}$, k_0 being negative. The values of the remaining free parameters are $\Delta = 30\lambda$ and $\beta = 0.995$ or $\gamma = 10$.

(cross-sectional) plane a one-photon field given by the FXW possesses the same strong localization at any time as the previously considered cylindrical field does in any transversal plane at the instant $t = 0$.

By making use of the historically first representative of localized waves – the so-called focus wave mode (FWM) [22]-[24] (see also Ref. [10] and reviews [11], [15] and references therein) one readily obtains an example of

the field that exhibits even much stronger than exponential localization. FWM is given by the scalar function

$$\Psi(\rho, z, t) = \frac{\exp\left[-\frac{\rho^2}{2l(a - i(z - \tau))}\right]}{a - i(z - \tau)} \exp\left[-\frac{i(z + \tau)}{2l}\right], \quad (10)$$

where again l is a wavelength-type characteristic length and the constant a controls the axial localization length. Since the FXW in the limit $\beta \rightarrow 1$ becomes a FWM [10], Fig.2 gives also an idea how a FWM looks like. Multiplying Eq.(10) by $\mathbf{m}$ to build the vector $\mathbf{Z}(\rho, z, \tau)$ and inserting the latter into Eq.(4) we obtain that in this example the photon localization in the waist plane is quadratically exponential (Gaussian falloff):

$$|\mathbf{Z}(\rho \rightarrow \infty, z = \tau)| \sim \exp(-\rho^2/2la), \quad (11)$$

$$|\mathbf{F}(\rho \rightarrow \infty, z = \tau)|^2 \sim \rho^6 \exp(-\rho^2/la). \quad (12)$$

In Eq.(12) only the highest-power term with respect to ρ is shown.

To start discussing our results let us ask first whether the wave functions considered are something extraordinary. The answer is: yes, they are indeed, since the browsing of various integral transform tables reveals rather few examples where both the real and imaginary part of a wave function and of its time derivative have simultaneously an exponential or stronger localization in conjunction with other requisite properties. Still, the list of proper wave functions with an extraordinary strong localization is not poor – in addition to an optically feasible version [25] of the FWM various new interesting solutions can be derived [26]. Yet, it could be argued that the well-known Gaussian beam pulse has the same quadratically exponential radial profile in the waist region. However, resorting to the family of the Gaussian beams (the Gauss-Laguerre and Gauss-Hermite beams, etc.) is irrelevant here. The reason is that all these beams are solutions of the wave equation only in the paraxial approximation not valid in the case of any significant localization of wide-band (pulsed) superpositions of the beams, whereas in fact, e.g., an exact solution corresponding to a lowest-order (axisymmetric) Gaussian beam has a weak power-law radial falloff in the waist region [27],[28].

The next possible objection to the physical significance of the results obtained might arise from the infinite total energy [11] of the waves given by Eqs. (6), (9), and (10). However, at any spatial location the wave function is square integrable with respect to time, thus the condition of the Paley-Wiener theorem has been satisfied. Moreover, physically feasible finite-energy versions of localized waves generally exhibit even better localization properties, although not persistently. A finite-energy version of the FXW, called the modified focused X wave (MFXW [11]), has the same exponential factor as in Eq.(9), which is multiplied by a fraction that allows to force the axial localization to follow an arbitrarily strong

power-law. The latter circumstance indicates that the strong lateral localization of the fields considered does not appear somehow at the expense of their axial localization. As a matter of fact, energy-normalization of a wave function depends on how many photons it describes. It is easy to see that our derivation and results hold for any number state with $N \geq 1$ and also for incoherent mixtures of such states (which is important for experimental studies). In contrast, the localization of coherent states is not governed by the Paley-Wiener theorem [6].

The final crucial question is, are our results not in contradiction with those of Ref. [6]? The answer is no, since in the case of the cylindrical waves the radial distance and temporal frequency are not directly Fourier-conjugated variables. In order to clarify this point, let us first take a closer look at the proof of the Paley-Wiener limit for three-dimensional isotropic localization. In Ref. [6] Eq.(24) for $\mathbf{Z}(\mathbf{r}, \tau)$ contains a superposition of spherically symmetric standing waves

$$\int_0^\infty dk h(kl) \frac{\sin kr}{r} e^{-ik\tau} \propto ir^{-1} \left[g\left(\frac{\tau+r}{l}\right) - g\left(\frac{\tau-r}{l}\right) \right], \quad (13)$$

where $h(kl)$ is the spectrum and $g(\cdot)$ is its Fourier image. The sine in Eq.(13) results from the imploding and exploding spherical wave constituents of the standing wave, like an odd one-dimensional standing wave arises from counterpropagating waves. While the asymptotic behavior of the function $g(\cdot)$ and hence of the function $\mathbf{Z}(\mathbf{r}, \tau)$ for large values of the radial distance r are generally restricted by the Paley-Wiener theorem, strictly at the instant $\tau = 0$ of maximal localization the integral is nothing but the sine transform for which the theorem does not apply. Indeed, the sine transform tables give examples of the resultant functions with arbitrarily abrupt falloff. However, it does not mean as if the localization restriction was lifted at the instant $\tau = 0$. The explanation is that according to Eq.(4) the energy wave function involves also the time derivative of $\mathbf{Z}(\mathbf{r}, \tau)$, but the sine transforms of two functions $h(kl)$ and $h(kl)k$ cannot simultaneously possess arbitrarily abrupt falloffs. In contrast, the time derivative of the wave function given by Eq.(6) or Eq.(5) has the same strong exponential falloff as the function itself, which persists for some (not too long) time, see Fig.1. By comparing Eqs. (5) and (13) we notice that while in Eq.(13) – as well as in its one-dimensional equivalent – the argument of the sine function is the product of the distance with the Fourier variable, in Eq.(5) the argument of the Bessel function is the product of the radial distance ρ with the radial wavenumber k_ρ the latter depending on the Fourier variable through the square-root expression with the constant parameter k_0 – the lower limit of the integration. As it follows also from Eqs. (7) and (8) the condition $k_0 \neq 0$ is crucial for obtaining the exponential falloff. Hence, in the case of the cylindrical waves considered by us, the apparent viola-

tion of the rules set by the Paley-Wiener theorem results from the specific complicated relation between the radial distance and the Fourier variable.

In conclusion, we have shown that for certain cylindrical N -photon states ($N = 1, 2, \dots$) the localization in lateral directions breaks the limit established in Ref. [6] for the case of uniform spherical wave functions. These results hold not only for photons but for number states of any particles.

The first author is thankful to Iwo Bialynicki-Birula for stimulating hints, discussions, and remarks. The research was supported by the Estonian Science Foundation.

-
- [1] L. D. Landau and R. Peierls, *Z. Phys.* **62**, 188 (1930).
 - [2] I. Bialynicki-Birula, in *Progress in Optics*, edited by E. Wolf (Elsevier, Amsterdam, 1996), Vol. XXXVI.
 - [3] A. L. Akhiezer and V. B. Berestetskii, *Quantum Electrodynamics* (Interscience, New York, 1965).
 - [4] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, Cambridge, 1995).
 - [5] M. Hawton, *Phys. Rev. A*, **59**, 3223 (1999).
 - [6] I. Bialynicki-Birula, *Phys. Rev. Lett.* **80**, 5247 (1998).
 - [7] O. Keller, *Phys. Rev. A*, **62**, 022111 (2000).
 - [8] O. Keller, *JOSA B*, **18**, 206 (2001).
 - [9] K. W. Chan, C. K. Law, and J. H. Eberly, *Phys. Rev. Lett.* **88**, 402 (2002).
 - [10] P. Saari and K. Reivelt, *Phys. Rev. E* **69**, 036612 (2004).
 - [11] I. Bieseris, M. Abdel-Rahman, A. Shaarawi, and A. Chatzipetros, *Progr. in Electrom. Res.* **19**, 1 (1998).
 - [12] J. Salo, J. Fagerholm, A. T. Friberg, and M. M. Salomaa, *Phys. Rev. E*, **62**, 4261 (2000).
 - [13] P. Saari, in *Time's Arrows, Quantum Measurement and Superluminal Behavior*, eds. D. Mugnai et al (CNR, Rome, 2001), p. 37 or arXiv: physics/0103054 (2002).
 - [14] E. Recami, M. Zamboni-Rached, K. Z. Nóbrega, C. A. Dartora, and H. E. Hernández, *IEEE J. Selected Topics in Quantum Electronics*, **9**, 59 (2003).
 - [15] K. Reivelt and P. Saari, arXiv: physics/0309079 (2003).
 - [16] P. Saari and K. Reivelt, *Phys. Rev. Lett.*, **79**, 4135 (1997).
 - [17] H. Sönajalg, M. Rätsep, and P. Saari, *Opt. Lett.* **22**, 310 (1997).
 - [18] D. Mugnai, A. Ranfagni, and R. Ruggeri, *Phys. Rev. Lett.* **84**, 4830 (2000).
 - [19] I. Alexeev, K. Y. Kim, and H. M. Milchberg, *Phys. Rev. Lett.* **88**, 073901 (2002).
 - [20] K. Reivelt and P. Saari, *Phys. Rev. E* **66**, 056611 (2002).
 - [21] R. Grunwald, V. Kebbel, U. Griebner, U. Neumann, A. Kummrow, M. Rini, E. T. J. Nibbering, M. Piché, G. Rousseau, and M. Fortin, *Phys. Rev. A* **67**, 063820 (2003).
 - [22] J. N. Brittingham, *J. Appl. Phys.* **54** 1179, (1983).
 - [23] A. Sezginer, *J. Appl. Phys.* **57**, 678 (1984).
 - [24] R. W. Ziolkowski, *J. Math. Phys.* **26**, 861 (1985).
 - [25] K. Reivelt and P. Saari, *Opt. Lett.*, **29**, 1176 (2004).
 - [26] A. P. Kiselev, *J. Phys. A: Math. Gen.*, **36**, L345 (2003).
 - [27] C. J. R. Sheppard and S. Saghafi, *JOSA A*, **16**, 1381 (1999).
 - [28] P. Saari, *Opt. Express* **8**, 590 (2001).